

DFTT 64/92
October 1992

VIOLATION OF THE BJORKEN SUM RULE IN QCD

Stefano Forte

*I.N.F.N., Sezione di Torino
via P. Giuria 1, I-10125 Torino, Italy*

ABSTRACT

We show that the Bjorken sum rule for the first moment of the polarized nucleon structure function g_1 is not necessarily satisfied in QCD if the axial anomaly is taken into account. We argue that nonperturbative QCD effects should lead to a violation of the sum rule of the order of the percentage isospin violation in current quark masses.

Submitted to: *Physics Letters B*

The Bjorken sum rule [1], which relates the isotriplet component of the first moment of the polarized nucleon structure function g_1 to the weak couplings G_A and G_V is usually considered a solid consequence of the standard model, and has been traditionally viewed as a benchmark of current universality and the quark interpretation. Even though this sum rule has never been tested directly, it is routinely used in the analysis and extraction of polarized structure function data.

Recent high-precision experimental determinations of first moments of polarized [2] (for a discussion see [3]) and unpolarized [4] (for a discussion see [5]) structure functions have lead to results which disagree with naive expectations based on partonic intuition, and have lead to reconsider assumptions, such as the isoscalarity of the antiquark sea in the nucleon, which, even though not a consequence of QCD, were always made in the analysis of experimental data. It is thus reasonable to ask, what is the status of the Bjorken sum rule in QCD? Two recent statements which express the common wisdom of, respectively, the experimental and theoretical communities in this respect are “today the Bjorken sum rule is regarded as a test of QCD”[6], and “the Bjorken sum rule presents a basic test of the current algebra”[7].

Here we shall show that, whereas the latter statement is correct, the former is not: it is possible to violate the Bjorken sum rule without violating QCD — nor its current algebra, which, due to the U(1) problem, is not the same as the quark current algebra [8]. In particular, due to a gluonic admixture to matrix elements of the singlet axial current, triggered by the axial anomaly, the Bjorken sum rule is only true provided one makes an assumption of isoscalarity of the gluon sea. This assumption, although plausible, need not be true, and actually, we shall argue that non-perturbative gluon configurations (like

instantons) should violate the assumption if the (light) current quark masses are unequal (as they are in the real world).

Let us first review the standard derivation of the Bjorken sum rule. Both current-algebra arguments [7] and the operator-product expansion [3] lead to equate the first moment of the polarized structure function g_1 with the (forward) matrix element of the electromagnetic axial current:

$$\int dx g_1(x) = C A_1 \quad (1)$$

$$A_1 s^\mu \equiv \langle N | \sum_i q_i^2 \bar{\psi}_i \gamma^\mu \gamma_5 \psi_i | N \rangle,$$

where s^μ is the spin four-vector of the given hadronic target N , $s_\mu s^\mu = -1$, and the sum runs over quark flavors with charge q_i . The coefficient C is computed perturbatively as $C = \frac{1}{2} \left(1 - \frac{\alpha_s}{\pi} + O(\alpha_s^2) \right)$. In order to extract a useful sum rule from Eq.(1) one takes an isotriplet matrix element, i.e., the proton-neutron difference

$$\int dx (g_1^p(x) - g_1^n(x)) = -s_\mu C \langle N | \sum_i q_i^2 \bar{\psi}_i \gamma^\mu \gamma_5 \psi_i | N \rangle^{I=1}, \quad (2)$$

then one uses current universality to express the isotriplet matrix element on the r.h.s. of Eq.(2) as the matrix element of the isotriplet part of the current, i.e.,

$$\int dx (g_1^p(x) - g_1^n(x)) = -s_\mu C \langle N | \left(\sum_i q_i^2 \bar{\psi}_i \gamma^\mu \gamma_5 \psi_i \right)^{I=1} | N \rangle \quad (3)$$

$$= C \frac{1}{3} \frac{G_A}{G_V}, \quad (4)$$

where the last equality — the Bjorken sum rule — which relates the matrix element of the isotriplet current to the weak couplings G_A , G_V , follows from straightforward isospin algebra.

Whereas the derivation of Eq.(1) is based on QCD current algebra and asymptotic freedom, and thus a failure of that equation would surely undermine QCD, in order to get to the Bjorken sum rule proper (4) one needs the extra step leading from Eq.(2) to Eq.(3). Even though this seems quite innocuous, the assumption on which is based taken at face value is obviously false. Indeed, whereas the quark currents form a $U_V(3) \times U_A(3)$ current algebra, at the hadronic level this is broken down to $U_V(3) \times SU_A(3)$ because of the axial anomaly which removes the singlet axial $U_A(1)$ symmetry [8]. This is manifested, for instance, by the fact that the matrix elements of the singlet current are scale dependent, while those of the nonsinglet one aren't. It follows that the singlet current's matrix elements are not in the same multiplet as those of the nonsinglet ones, thus we do not know their transformation properties under isospin.

We can see this more explicitly by introducing a decomposition [3], [9] of the form factor G_A of the forward nucleon matrix element of the isosinglet axial current in terms of its quark components Δq_i and gluon component ΔG :

$$\begin{aligned}
G_A s^\mu &\equiv \langle N | \sum_i \bar{\psi}_i \gamma^\mu \gamma_5 \psi_i | N \rangle \\
G_A &= \sum_i (\Delta q_i - \Delta G) \\
\Delta G &= (\Omega + \Delta g),
\end{aligned} \tag{5}$$

where Δg and Ω are, respectively, the contributions from perturbative and non-perturbative gluonic configurations to ΔG . Despite some controversies on the way the gluonic contribution could be experimentally singled out, it is now clear [10] that the decomposition (5) can be established in a gauge-invariant way both perturbatively in the QCD parton model [3] and nonperturbatively at the operator level [9]. The simplest way of introducing it is to observe [9] that the matrix element of the singlet axial charge $Q_5 \equiv \int d^3x \sum_i \bar{\psi}_i \gamma^0 \gamma_5 \psi_i$

is, in general, the sum of the canonical quark helicity operator Q_5^q and a gluonic operator Q_5^g , whose nucleon matrix elements are respectively identified with Δq and ΔG . Whereas Q_5^q is conserved, Q_5^g is not and the gluonic contribution is thus responsible for the anomalous divergence of the axial current[11]. The separation of the gluonic contribution in a perturbative and a nonperturbative portion is discussed in detail in Ref.[9], and can be understood [10] observing that the (perturbative) gluon distribution provides a contribution Δq to ΔG which evolves at two loops, while nonperturbative field configurations (such as instantons) provide a scale-invariant contribution.

Using Eq.(5) in Eq.(2), neglecting QCD corrections, and assuming strange and heavier flavor distributions to be isosinglet to good approximation (we shall come back on this later) we get

$$\begin{aligned} \int dx (g_1^p(x) - g_1^n(x)) &= \\ &= \frac{1}{2} \left[-s_\mu \langle N | \left(\frac{4}{9} \bar{u} \gamma^\mu \gamma_5 u + \frac{1}{9} \bar{d} \gamma^\mu \gamma_5 d \right) | N \rangle^{I=1} - \frac{5}{9} \langle N | \Delta G | N \rangle^{I=1} \right]. \end{aligned} \quad (6)$$

The (non-anomalous) quark contributions Δq to G_A do belong in the full $U(3) \times U(3)$ multiplet, thus we can use isospin symmetry to relate $\Delta u^p = \Delta d^n$ and so forth (where the superscript denotes the state in which the matrix element is taken) to get

$$\begin{aligned} \int dx (g_1^p(x) - g_1^n(x)) &= \frac{1}{2} \left[\frac{1}{3} (\Delta u^p - \Delta d^p) - \frac{5}{9} (\Delta G^p - \Delta G^n) \right] \\ &= \frac{1}{2} \langle N | \left(\sum_i q_i^2 \bar{\psi}_i \gamma^\mu \gamma_5 \psi_i \right)^{I=1} | N \rangle - \frac{5}{18} (\Delta G^p - \Delta G^n) \\ &= C \left[\frac{1}{3} \left(\frac{G_A}{G_V} \right) - \frac{5}{9} (\Delta G^p - \Delta G^n) \right]. \end{aligned} \quad (7)$$

It follows that the Bjorken sum rule is corrected at the crucial step (3) by the last term on the r.h.s. of Eq.(7), and it is true only upon the assumption that this term vanishes.

We have thus shown explicitly that the Bjorken sum rule follows from QCD only upon the assumption that the gluonic component ΔG is isosinglet. Because there are now reasons to believe that ΔG may be actually rather large (i.e., comparable to Δq_i) [3], it is important to see whether we should expect indeed ΔG to be isosinglet or not. It should be realized that since ΔG receives in general both perturbative and nonperturbative contributions, there is no *a priori* reason why it should be isosinglet. Rather, this is a phenomenological assumption on the same footing as that according to which the nucleon sea should be SU(2) symmetric, which has recently been disproved experimentally [4],[5].

Indeed, isospin violations of order 5% in the sea parton distributions are suggested [5] by recent data [4], and isospin violation of, for example, the nucleon radii up to about 0.2 Fm are compatible with current electron-nucleus scattering data [12]. Due to the numerical coefficients in Eq.(7) assuming [3],[9] $|\Delta G| \sim \frac{1}{N_f} \Delta q$, this leads to expect violation of the Bjorken sum rule of order 2 – 3% just from isospin violation in the gluon distributions. If ΔG receives a sizable nonperturbative contribution, however, even larger violations of the Bjorken sum rule may appear.

To understand this, it is convenient to consider the matrix element of the divergence of the (singlet) axial current, which is given by the anomaly equation [11]:

$$G_A s \cdot (p' - p) \equiv \langle N | \partial_\mu j_5^\mu | N \rangle = \langle N | \sum_i 2m_i \bar{\psi} i \gamma_5 \psi + \frac{N_f}{8\pi^2} g^2 \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} | N \rangle, \quad (8)$$

where $(p' - p)$ is the momentum transfer and m_i is the current quark mass for the i -th flavor. In order to avoid possible sources of confusion, it should be pointed out rightaway that the naive identification of the two terms on the r.h.s. of Eq.(8) with the quark and gluon contributions Δq and ΔG is incorrect [13]. Next, it is convenient to define [14][15], [13](see

also Ref.[16]) one-meson reducible and one-meson irreducible contributions to the matrix element on the r.h.s. of Eq.(8). The former are essentially given [14] by diagrams where the mass term or the gluon condensate couples to a pion (or generally a pseudoscalar meson) with the appropriate quantum numbers, which then couples to the nucleon. A typical contribution to the latter is that where the mass term couples directly to one of the nucleon's quarks.

It is easy to see [14] that the one-particle reducible contribution to the matrix element of the mass term on the r.h.s. of Eq.(8) displays large isospin violation, because the singlet mass terms couples to the triplet meson (the pion) with amplitude [14]

$$\langle 0 | \left(\sum_i 2m_i \bar{\psi} i \gamma_5 \psi \right) | \pi^0 \rangle = \frac{m_u - m_d}{m_u + m_d} f_\pi M_\pi^2, \quad (9)$$

hence, neglecting the η contribution to the reducible portion of the matrix element of Eq.(8), and retaining only the pion one, leads immediately to isospin violation of order $\frac{m_u - m_d}{m_u + m_d}$. Because $\frac{m_d}{m_u} \approx 2$ [17] this is a very large violation. However, as brilliantly shown in Ref.[14], the Sutherland-Veltman theorem implies the vanishing of the one-pion matrix element of the *full* r.h.s. of Eq.(8), which in turn implies that the one-meson reducible matrix element of the anomaly must display an isospin violation equal and opposite to that of the mass term, such that the overall reducible contribution to the matrix element of $\partial_\mu j_5^\mu$ is isospin conserving. This is borne out by explicit effective lagrangian computations [15]. This justifies the neglect of heavy quark contributions to Eq.(6), since the isospin violation of these is entirely compensated by that of the anomaly [14].

What about the irreducible contributions? Despite the explicit isospin violation from the (current) quark masses, the matrix element of the mass term in a quark state with helicity λ is perfectly isospin invariant:

$$\langle q | m \bar{\psi} i \gamma_5 \psi | q \rangle = 2\lambda(p' - p) \cdot s \quad (10)$$

because its mass dependence cancels. Notice that this implies that the mass term cannot be neglected even in the chiral limit; if this term is incorrectly dropped from the Ward identities derived from Eq.(8) one reaches [18] the absurd conclusion that the helicity of a free quark vanishes in the chiral limit. Eq.(10) leads thus to expect that no isospin violation should arise from the irreducible matrix elements of the mass term.

It is interesting to consider instead irreducible contributions to the matrix element of the anomaly from instanton-like field configurations. These have been computed in Ref.[19] for a simplified model of a single massless quark, in the one-flavor case. In that Ref., it was shown that the matrix element of the anomaly receives a contribution from instanton-induced couplings which can be identified with a contribution to Ω Eq.(5), and which exactly cancels the bare quark helicity, $\Delta q = -\Delta G$. This suggests that the nonperturbative gluonic contribution Ω to the nucleon matrix element of the axial current Eq.(5) may also be large and anticorrelated to Δq , thus explaining the surprising experimental discovery [2] that this matrix element vanishes (within experimental uncertainties).

Such contribution arises from a direct coupling of the quark line to instantons and is thus irreducible according to the above classification. Now the effective instanton-induced chirality flipping interaction which generates it arises due to fermionic zero modes in the instanton field, and is thus suppressed by $\frac{1}{m_q}$ if the quark is given a mass m_q . This leads one to expect that in the many-flavor case the corresponding contribution (which in the mean-field approximation is identical to the one-flavor one [19]) should display isospin violations of order $\frac{\Omega^p}{\Omega^n} \sim \frac{m_d}{m_u}$.

In order to get a feeling for the size of the expected violation of the sum rule, we can very crudely assume that Δq_i take the values obtained from the quark model and the

Ellis-Jaffe sum rule (see [3]), screened by Ω so that $A_1 = 0$, while $\frac{\Omega^p}{\Omega^n} \sim \frac{m_d}{m_u}$. This leads to estimate the correction in Eq.(7) to be $\frac{5}{18}(\Delta G^p - \Delta G^n) \sim 0.1$, which is roughly one-half of the Bjorken value $\frac{1}{6} \frac{G_A}{G_V}$. Otherwise stated, the extra term on the r.h.s. of the Bjorken sum rule would thus be estimated to reduce the Bjorken value by roughly a factor 2.

How would such a dramatic violation of the Bjorken sum rule affect the analysis of data on the polarized proton structure function which has been presented by the EMC collaboration [2], and which has stirred so much attention in the physics community? A minute's reflection reveals that it wouldn't affect it at all. In order to extract from the data, which provide a value of the first moment of the proton structure function $\int dx g_1(x)$, a value for the singlet form factor G_A Eq.(5), which is of more direct physical significance, one needs to use Eq.(4), expressed as

$$\Delta u - \Delta d = \frac{G_A}{G_V}, \quad (11)$$

together with an analogous relation for the SU(3) octet current. Now Eq.(11) (and Eq.(4)) are perfectly correct, regardless of whether the assumption leading from Eq.(2) to Eq.(3) fails thereby leading to violation of the Bjorken sum rule. The existence of such a violation would only imply that, due to the presence of the gluonic contribution in Eqs.(5),(7), knowledge of the proton value of both the first moment A_1 Eq.(1) and the singlet form factor G_A Eq.twocomp is insufficient to determine the respective neutron values.

Rather than providing evidence against QCD, a violation of the Bjorken sum rule would thus provide evidence in favor of the fact that indeed the matrix element of the singlet axial current receives an extra gluonic contribution, which is necessarily absent in the matrix elements of the nonsinglet currents, since this is the only possible source of

isospin violation in the proton-neutron difference of structure functions. Even though it will not provide a test of QCD, contrary to widespread belief, the forthcoming experimental test of the Bjorken sum rule will provide us with extremely valuable information on the nonperturbative structure of this theory.

Acknowledgements: I thank M. Anselmino, M. Arneodo, and E. Predazzi for discussions.

REFERENCES

References

- [1] J. D. Bjorken, *Phys. Rev* **148**, 1467 (1966)
- [2] J. Ashman *et al.*, *Nucl. Phys.* **B328**, 1 (1990)
- [3] G. Altarelli, in ‘The Challenging Questions’, Proceedings of the 1989 Erice school, A. Zichichi, ed. (Plenum, NY, 1990)
- [4] P. Amaudruz *et al.*, *Phys. Rev. Lett.* **66**, 560 (1991)
- [5] S. Forte, Torino preprint DFTT 18/92 (1992), *Phys. Rev. D*, in press
- [6] C. Y. Prescott, SLAC Preprint SLAC-PUB-5604 (1991)
- [7] S. D. Drell, SLAC Preprint SLAC-PUB-5720 (1992)
- [8] See e.g. S. Coleman, ‘Aspects of Symmetry’ (Cambridge, 1985)
- [9] S. Forte, *Phys. Lett.* **B224**, 189 (1989); *Nucl. Phys.* **B331**, (1990)
- [10] See G. Altarelli, CERN preprint CERN-TH.6340/91 (December 1991)
- [11] See R. Jackiw, in S. B. Treiman, R. Jackiw, B. Zumino and E. Witten, ‘Current Algebra and Anomalies’ (World Scientific, Singapore, 1985)
- [12] T. W. Donnelly, J. Dubach and I. Sick, *Nucl. Phys.* **A503**, 589 (1989)
- [13] G. M. Shore and G. Veneziano *Phys. Lett.* **B244**, 75 (1990); *Nucl. Phys.* **B381**, 3 (1992)
- [14] D. G. Gross, S. Treiman and F. Wilczek, *Phys. Rev.* **D19**, 2188 (1977)
- [15] G. Veneziano, *Mod. Phys. Lett.* **A4**, 1605 (1989)
- [16] K. F. Liu, *Phys. Lett.* **B281**, 141 (1992)
- [17] S. Weinberg, in ‘A Festschrift for I. I. Rabi, L. Motz, ed. (New York Academy of Sciences, New York, 1977)
- [18] R. L. Jaffe and A. Manohar, *Nucl. Phys.* **B337**, 509 (1990)
- [19] S. Forte and E. V. Shuryak, *Nucl. Phys.* **B357**, 153 (1991)